

Examples for Souslin Forcing*

Haim Judah

Dept. of Mathematics and Computer Science
Bar-Ilan University
52900 Ramat-Gan, Israel

Andrzej Rosłanowski

Dept. of Mathematics and Computer Science
Bar-Ilan University
52900 Ramat-Gan, Israel

and

Mathematical Institute of Wroclaw University
50384 Wroclaw, Poland

Saharon Shelah

Institute of Mathematics
The Hebrew University of Jerusalem
Jerusalem, Israel

and

Department of Mathematics
Rutgers University Rutgers, USA

February 1, 2008

*The authors would like to thank Israel Academy of Sciences BRF and MSRI for partial support.

1 Introduction

In this paper we continue with study of forcing notions having a simple definition. We began this study in [JS1] and [JS2]. In [BJ] we gave more results about Souslin forcing notions and in this paper we will give some examples of Souslin forcing notions answering a question of [JS1] and a question of H.Woodin.

A forcing notion $\mathbf{P}$ is Souslin if $\mathbf{P} \subseteq \mathcal{R}$ is a Σ_1^1 -set, $\{(p, q) : p \leq_{\mathbf{P}} q\}$ is a Σ_1^1 -set and $\{(p, q) : p \text{ is incompatible with } q\}$ is a Σ_1^1 -set.

More information on Souslin forcing notion can be found in [JS1]. A related work is [BJ]. In [JS1] we prove that if $\mathbf{P}$ is Souslin ccc and $\mathbf{Q}$ is any forcing notion then $\mathbf{V}^{\mathbf{Q}} \models \text{“}\mathbf{P} \text{ satisfies ccc”}$. A natural question was: does “ $\mathbf{P}$ is Souslin ccc” imply “ $\mathbf{P}$ has Knaster property”. Recall that $\mathbf{P}$ satisfies Knaster property if and only if

$$(\forall A \in [\mathbf{P}]^{\omega_1})(\exists B \in [A]^{\omega_1})(\forall p, q \in B)(p \text{ is compatible with } q).$$

In the second section we will give a model where there is a ccc Souslin forcing which does not satisfy the Knaster condition. Recall that under the assumption of $\mathbf{MA}$ every ccc notion of forcing has Knaster property.

Many simple forcing notions $\mathbf{P}$ satisfy the following condition:

$$\Vdash_{\mathbf{P}} \text{“}\hat{\mathbf{P}} \text{ is } \sigma\text{-centered”}.$$

This property is connected with the homogeneity of the forcing notion. The example of a totally nonhomogeneous Souslin forcing will be constructed in the third section.

In the next section we present a model where there is a σ -linked not σ -centered Souslin forcing such that all its small subsets are σ -centered but Martin Axiom fails for this order.

In section 5 we will give an example of a σ -centered Souslin forcing notion and a model of the negation of $\mathbf{CH}$ in which the union of less than continuum meager subsets of $\mathcal{R}$ is meager but Martin Axiom fails for this notion of forcing.

In the last session of the MSRI Workshop on the continuum (October 1989) H.Woodin asked if “ $\mathbf{P}$ has a simple definition and does not satisfy ccc” implies that there exists a perfect set of mutually incompatible conditions. Clearly Mathias forcing satisfies such a requirement. In section 6 we

will find a Souslin forcing which is Proper but not ccc that does not contain a perfect set of mutually incompatible conditions.

The last section will show that ccc Σ_2^1 -notions of forcing may not be indestructible ccc.

Our notation is standard and derived from [Je]. There is one exception, however. We write $p \leq q$ to say that q is a stronger condition than p .

2 On the Knaster condition

In this section we will build a Souslin forcing satisfying the countable chain condition but which fails the Knaster condition.

Fix a sequence $\langle \sigma_i : i \in \omega \rangle$ of functions from ω into ω such that

(*) if $N < \omega$, $\phi_i : N \longrightarrow \omega$ (for $i < N$) then there are distinct $n_0, \dots, n_{N-1}$ such that

$$(\forall i, j_0, j_1 < N)(\phi_i(j_0) = j_1 \Rightarrow \sigma_i(n_{j_0}) = n_{j_1}).$$

Note that there exists a sequence $\langle \sigma_i : i \in \omega \rangle$ satisfying (*):

Suppose we have defined $\sigma_i|_{m_0} : m_0 \longrightarrow m_0$ for $i < m_0$. We want to ensure (*) for $n_0 + 1, \phi_i$ ($i \leq n_0$). Define $\sigma_i(m_0 + j_0) = m_0 + \phi_i(j_0)$ for $i, j_0 \leq n_0$. Take large m_1 and extend all σ_i ($i \leq n_0$) on m_1 in such a way that $\text{rng}(\sigma_i) \subseteq m_1$.

Next we define functions $f_i : \omega^\omega \longrightarrow \omega^\omega$ for $i \in \omega$ by

$$f_i(x)(k) = \begin{cases} x(k) & \text{if } k < i \\ \sigma_i(x(k)) & \text{otherwise} \end{cases}$$

Clearly all functions f_i are continuous. Put $F(x) = \{f_i(x) : i \in \omega\}$ for $x \in \omega^\omega$.

Lemma 2.1 *Suppose that $x_\alpha, y_\alpha \in \omega^\omega$ are such that there is no repetition in $\{x_\alpha, y_\alpha : \alpha \in \omega_1\}$. Then there exists $A \in [\omega_1]^{\omega_1}$ such that*

$$(\forall \alpha, \beta \in A)(\alpha < \beta \Rightarrow x_\alpha \notin F(y_\beta)).$$

PROOF For $\alpha < \omega_1$ let $n_\alpha = \min\{n : x_\alpha(n) \neq y_\alpha(n)\}$. We find $n \in \omega$, and $s, t \in \omega^n$ such that the set $A_0 = \{\alpha < \omega_1 : n = n_\alpha + 1 \text{ \& } x_\alpha|n = s \text{ \& } y_\alpha|n = t\}$ is stationary in ω_1 . Clearly $s \neq t$ and $s|(n-1) = t|(n-1)$. Thus $\alpha, \beta \in A_0$ and $x_\alpha \in F(y_\beta)$ imply $x_\alpha \in \{f_i(y_\beta) : i \leq n\}$. Consequently the set $\{\alpha \in A_0 \cap \beta : x_\alpha \in F(y_\beta)\}$ is finite for each $\beta \in A_0$.

We define the regressive function $\psi : A_0 \longrightarrow \omega_1$ by $\psi(\beta) = \max\{\alpha \in A_0 \cap \beta : x_\alpha \in F(y_\beta)\}$ (with the convention that $\max \emptyset = 0$). By Fodor's lemma there are $\gamma < \omega_1$ and a stationary set $A_1 \subseteq A_0$ such that $\psi(\beta) = \gamma$ for all $\beta \in A_1$. Put $A = A_1 \setminus (\gamma + 1)$. Now, if $\alpha, \beta \in A$, $\alpha < \beta$ then $\psi(\beta) < \alpha$ and hence $x_\alpha \notin F(y_\beta)$. ■

Lemma 2.2 *Suppose that $\{W_\alpha : \alpha < \omega_1\}$ is a family of disjoint finite subsets of ω^ω . Then there exist $\beta < \omega_1$ and an infinite set $A \subseteq \beta$ such that*

$$(\forall \alpha \in A)(\forall x \in W_\alpha)(\forall y \in W_\beta)(x \notin F(y)).$$

PROOF We may assume that all sets W_α are of the same cardinality, say $|W| = n$ for $\alpha < \omega_1$. For $\alpha = \lambda + k$, where $\lambda < \omega_1$ is a limit ordinal and $k \in \omega$ we define $X_\alpha = W_{\lambda+2k}$ and $Y_\alpha = W_{\lambda+2k+1}$. Let $X_\alpha = \{x_i^\alpha : i < n\}$, $Y_\alpha = \{y_i^\alpha : i < n\}$. Choose by the induction on $l = l_1 \cdot n + l_2 < n^2$, $l_1, l_2 < n$ uncountable sets $A_l \subseteq \omega_1$ satisfying

- $A_{l+1} \subseteq A_l$ and
- if $l = l_1 \cdot n + l_2$, $l_1, l_2 < n$, $\alpha, \beta \in A_l$ and $\alpha < \beta$ then $x_{l_1}^\alpha \notin F(y_{l_2}^\beta)$.

Since there is no repetition in $\{x_{l_1}^\alpha, y_{l_2}^\alpha : \alpha \in A_{l-1}\}$ we may apply lemma 2.1 to get A_l from A_{l-1} .

Consider A_{n^2-1} . Choose $\beta_0 \in A_{n^2-1}$ such that the set

$$A = \{\lambda + 2k < \beta_0 : \lambda + k \in A_{n^2-1} \text{ \& } k \in \omega \text{ \& } \lambda \text{ is a limit ordinal}\}$$

is infinite. Let $\beta_0 = \lambda_0 + k_0$ where $k_0 \in \omega$ and λ_0 is limit. Put $\beta = \lambda_0 + 2k_0 + 1$. Since $\beta_0 < \beta$ we have $A \subseteq \beta$. Suppose $\alpha = \lambda + 2k \in A$. Let $x \in W_\alpha$, $y \in W_\beta$. Then $\lambda + k \in A_{n^2-1}$, $W_\alpha = X_{\lambda+k}$ and $W_\beta = Y_{\lambda_0+k_0} = Y_{\beta_0}$. Thus for some $l_1, l_2 < n$ we have $x = x_{l_1}^{\lambda+k}$ and $y = y_{l_2}^{\beta_0}$. Since $\lambda + k, \beta_0 \in A_{l_1 \cdot n + l_2}$, $\lambda + k < \beta_0$ we get $x \notin F(y)$. The lemma is proved. ■

Let relations R_i on $\omega^{<\omega}$ be defined by

sR_it if and only if
 $i < |s| = |t|$, $s \upharpoonright i = t \upharpoonright i$ and $(\forall l \in [i, |s|))(s(l) = \sigma_i(t(l)))$.

Note that if $x, y \in \omega^\omega$ are such that $(\forall n > i)(x \upharpoonright n R_i y \upharpoonright n)$ then $x = f_i(y)$.

We define the following forcing notion $\mathbf{Q}$. A member q of $\mathbf{Q}$ is a finite function such that:

- $\alpha)$ $\text{dom}(q) \in [\omega_1]^{<\omega}$, $\text{rng}(q) \subseteq \omega^{<\omega}$,
- $\beta)$ $(\forall \alpha, \beta \in \text{dom}(q))(\alpha \neq \beta \Rightarrow q(\alpha) \neq q(\beta))$,
- $\gamma)$ there is $n(q) \in \omega$ such that $q(\alpha) \in \omega^{n(q)}$ for all $\alpha \in \text{dom}(q)$.

The order is defined as follows:

$q \leq p$ if and only if

- 1. $\text{dom}(q) \subseteq \text{dom}(p)$ and
- 2. $(\forall \alpha \in \text{dom}(q))(q(\alpha) \subseteq p(\alpha))$ and
- 3. if $\alpha, \beta \in \text{dom}(q)$, $\alpha < \beta$, $i < n(q)$ and $q(\alpha) R_i q(\beta)$ then $p(\alpha) R_i p(\beta)$.

Lemma 2.3 $\mathbf{Q}$ satisfies ccc.

PROOF Suppose $\{q_\alpha : \alpha < \omega_1\} \subseteq \mathbf{Q}$. We find $\gamma < \omega_1$ and $A \in [\omega_1]^{\omega_1}$ such that for each $\alpha, \beta \in A$, $\alpha < \beta$ we have

- $n(q_\alpha) = n(q_\beta)$,
- $\text{dom}(q_\alpha) \cap \gamma = \text{dom}(q_\beta) \cap \gamma$, $(\text{dom}(q_\alpha) \setminus \gamma) \cap (\text{dom}(q_\beta) \setminus \gamma) = \emptyset$,
- $q_\alpha \upharpoonright (\text{dom}(q_\alpha) \cap \gamma) = q_\beta \upharpoonright (\text{dom}(q_\beta) \cap \gamma)$.

Suppose $\alpha, \beta \in A$. Clearly $\bar{q} = q_\alpha \cup q_\beta$ is a function. The only problem is that there may exist $\gamma_0 \in \text{dom}(q_\alpha)$ and $\gamma_1 \in \text{dom}(q_\beta)$ such that $q_\alpha(\gamma_0) = q_\beta(\gamma_1)$. Therefore to get a condition above both q_α and q_β we have to extend all $\bar{q}(\gamma)$. Let $\text{dom}(\bar{q}) = \{\gamma_j : j < N\}$ be an increasing enumeration. For $i < n(q_\alpha)$ choose $\phi_i : N \longrightarrow \omega$ such that

if $j_1 < j_0 < N$ and $\bar{q}(\gamma_{j_1})R_i\bar{q}(\gamma_{j_0})$ and either $\gamma_{j_0}, \gamma_{j_1} \in \text{dom}(q_\alpha)$
or $\gamma_{j_0}, \gamma_{j_1} \in \text{dom}(q_\beta)$
then $\phi_i(j_0) = j_1$.

Note that $\bar{q}(\gamma_{j_1})R_i\bar{q}(\gamma_{j_0}), \bar{q}(\gamma_{j_2})R_i\bar{q}(\gamma_{j_0})$ imply $\bar{q}(\gamma_{j_1}) = \bar{q}(\gamma_{j_2})$. Hence if $j_2 < j_1 < j_0$ are as above then $\gamma_{j_2} \geq \gamma$ and consequently only one pair (j_1, j_0) or (j_2, j_0) will be considered in the definition of ϕ_i . Apply condition $(*)$ to find distinct $n_0, \dots, n_{N-1}$ such that

$$(\forall i < n)(q_\alpha)(\forall j_0, j_1 < N)(\phi_i(j_0) = j_1 \Rightarrow \sigma_i(n_{j_0}) = n_{j_1}).$$

Put $q(\gamma_j) = \bar{q}(\gamma_j) \hat{\ } n_j$ for $j < N$. Clearly $q \in \mathbf{Q}$. Suppose $\gamma_{j_1}, \gamma_{j_0} \in \text{dom}(q_\alpha)$, $j_1 < j_0$ and $q_\alpha(\gamma_{j_1})R_iq_\alpha(\gamma_{j_0})$ for some $i < n(q_\alpha)$. Then $\phi_i(j_0) = j_1$ and hence $\sigma_i(n_{j_0}) = n_{j_1}$. Thus $q(\gamma_{j_1})R_iq(\gamma_{j_0})$. It shows that $q_\alpha \leq q$. Similarly $q_\beta \leq q$. Thus we have proved that $\mathbf{Q}$ satisfies Knaster condition. ■

Let $G \subseteq \mathbf{Q}$ be generic over $\mathbf{V}$. In $\mathbf{V}[G]$ we define $x_\alpha^G = \bigcup \{q(\alpha) : q \in G \ \& \ \alpha \in \text{dom}(q)\}$ for $\alpha < \omega_1$. Obviously each x_α^G is a sequence of integers. As in the proof of lemma 2.3 we can show that for each $q \in \mathbf{Q}$ there is $p \geq q$ such that $n(p) = n(q) + 1$. Consequently $x_\alpha^G \in \omega^\omega$ for every $\alpha < \omega_1$. Moreover $x_\alpha^G \neq x_\beta^G$ for $\alpha < \beta < \omega_1$ (recall that $q(\alpha) \neq q(\beta)$ for distinct $\alpha, \beta \in \text{dom}(q)$).

Note that if $\alpha < \beta$, $\alpha, \beta \in \text{dom}(q)$, $q \in \mathbf{Q}$ and $i < n(q)$ then

$$\begin{aligned} q(\alpha)R_iq(\beta) &\text{ implies } q \Vdash \dot{x}_\alpha = f_i(\dot{x}_\beta) \text{ and} \\ \neg q(\alpha)R_iq(\beta) &\text{ implies } q \Vdash \dot{x}_\alpha \neq f_i(\dot{x}_\beta). \end{aligned}$$

Lemma 2.4 *Suppose $G \subseteq \mathbf{Q}$ is generic over $\mathbf{V}$. Then*

$$\mathbf{V}[G] \models (\forall A \in [\omega_1]^{\omega_1})(\exists \alpha, \beta \in A)(\alpha < \beta \ \& \ x_\alpha^G \in F(x_\beta^G)).$$

PROOF Let $\dot{A}$ be a $\mathbf{Q}$ -name for an uncountable subset of ω_1 . Given $p \in \mathbf{Q}$. Find $A_0 \in [\omega_1]^{\omega_1}$ and $q_\alpha \geq p$ for $\alpha \in A_0$ such that $\alpha \in \text{dom}(q_\alpha)$ and $q_\alpha \Vdash \alpha \in \dot{A}$. We may assume that for each $\alpha, \beta \in A_0$ we have $n = n(q_\alpha) = n(q_\beta)$ and $q_\alpha(\alpha) = q_\beta(\beta)$. Now we repeat the procedure of lemma 2.3 with one small change. We choose suitable $A_1 \in [A_0]^{\omega_1}$, $\gamma < \omega_1$ and we take $\alpha, \beta \in A_1$, $\alpha < \beta$. Defining integers $n_0, \dots, n_{N-1}$ we consider functions $\phi_i : N \longrightarrow \omega$ (for $i < n$) as in 2.3 and a function $\phi_n : N \longrightarrow \omega$ such that $\phi_n(k) = l$, where $\alpha = \gamma_l$, $\beta = \gamma_k$. Then we get a condition $q \in \mathbf{Q}$ above both q_α and q_β and

such that $\sigma_n(q(\beta)(n)) = q(\alpha)(n)$. Since $q(\beta)|n = q(\alpha)|n$ and $n(q) = n + 1$ we have $q(\alpha)R_n q(\beta)$ and consequently $q \Vdash \dot{x}_\alpha = f_n(\dot{x}_\beta)$. Since $q \geq q_\alpha, q_\beta$ we get $q \Vdash “\alpha, \beta \in \dot{A} \ \& \ \dot{x}_\alpha \in F(\dot{x}_\beta)”$. ■

Fix a Borel isomorphism $(\pi_0, \pi_1, \pi_2) : \omega^\omega \longrightarrow (\omega^\omega)^\omega \times 2^\omega \times \omega \times \omega^\omega$. Thus if $x \in \omega^\omega$ then $\pi_1(x)$ is a relation on ω and $\pi_0(x)$ is a sequence of reals. Let Γ consists of all reals $x \in \omega^\omega$ such that

1. $(\forall n \neq m)(\pi_0(n) \neq \pi_0(m))$
2. $\pi_1(x)$ is a linear order on ω
3. $\pi_2(x) \in A_x = \{\pi_0(n) : n \in \omega\}$ and it is the $\pi_1(x)$ -last element of A_x .

Note that in 3 we think of $\pi_1(x)$ as an order on A_x . We define relations $<_\Gamma$ and $\equiv_\Gamma$ on Γ by

$x <_\Gamma y$ if and only if
 A_x is a proper $\pi_1(y)$ -initial segment of A_y and $\pi_1(y)|_{A_x} = \pi_1(x)$.

$x \equiv_\Gamma y$ if and only if
 $A_x = A_y$ and $\pi_1(y) = \pi_1(x)$ (we treat $\pi_1(x), \pi_1(y)$ as orders on A_x, A_y , respectively).

Clearly Γ is a Borel subset of ω^ω , $<_\Gamma$ is a Borel transitive relation on Γ and $\equiv_\Gamma$ is a Borel equivalence relation on Γ .

Now we define a forcing notion $\mathbf{P}_1$. Conditions in $\mathbf{P}_1$ are finite subsets p of Γ such that

if $x, y \in p$, $x <_\Gamma y$ then $\pi_2(x) \notin F(\pi_2(y))$.

$\mathbf{P}_1$ is ordered by the inclusion.

Lemma 2.5 $\mathbf{P}_1$ is a ccc Souslin forcing.

PROOF $\mathbf{P}_1$ is Souslin since it can be easily coded as a Borel subset of ω^ω in such a way that the order is Borel too. We have to show that $\mathbf{P}_1$ satisfies the countable chain condition. First let us note some properties of the incompatibility in $\mathbf{P}_1$. Suppose $p, q \in \mathbf{P}_1$ are incompatible. Clearly $p \setminus q$ and $q \setminus p$ are incompatible. If $x \in p$ and $x \equiv_\Gamma x'$ then $(p \setminus \{x\}) \cup \{x'\}$ and q are incompatible.

Suppose now that $\{p_\alpha : \alpha < \omega_1\}$ is an antichain in $\mathbf{P}_1$. By the Δ -lemma and by the above remarks we may assume that

- 1) $p_\alpha \cap p_\beta = \emptyset$ for $\alpha < \beta < \omega_1$
- 2) if $x, x' \in \bigcup_{\alpha < \omega_1} p_\alpha$, $x \neq x'$ then $x \not\equiv_\Gamma x'$.

Note that if $p \in \mathbf{P}_1$ then the set $\{[y]_{\equiv_\Gamma} : y \in \Gamma \text{ \& } (\exists x \in p)(y <_\Gamma x)\}$ is countable. Hence, due to 2, we may assume that

- 3) $(\forall \alpha < \beta < \omega_1)(\forall x \in p_\alpha)(\forall y \in p_\beta)(\neg y <_\Gamma x)$

CLAIM: Let $d \in [\omega^\omega]^{<\omega}$. Then $d = \{\pi_2(x) : x \in p_\alpha\}$ for at most countably many $\alpha < \omega_1$.

Indeed, assume not. Then we find $\beta < \omega_1$ such that $\{\pi_2(x) : x \in p_\beta\} = d$ and the set $B = \{\alpha < \beta : \{\pi_2(x) : x \in p_\alpha\} = d\}$ is infinite. Note that if $x', x'' <_\Gamma x$ and $\pi_2(x') = \pi_2(x'')$ then $x' \equiv_\Gamma x''$. Hence if $x \in p_\beta$ then for at most $|d|$ elements x' of $\bigcup_{\alpha \in A} p_\alpha$ we have $x' <_\Gamma x$. Thus we find $\alpha \in A$ such that $(\forall x' \in p_\alpha)(\forall x \in p_\beta)(\neg x' <_\Gamma x)$. It follows from 3 that

$$(\forall x \in p_\beta)(\forall x' \in p_\alpha)(\neg x <_\Gamma x')$$

and hence conditions p_α and p_β are compatible – a contradiction.

Let $d_\alpha = \{\pi_2(x) : x \in p_\alpha\}$. By the above claim we may assume that $d_\alpha \neq d_\beta$ for all $\alpha < \beta < \omega_1$. Applying the Δ -lemma we may assume that

- 4) $\{d_\alpha : \alpha < \omega_1\}$ forms a Δ -system with the root d .

Since the set $\bigcup_{w \in d} F(w)$ is countable w.l.o.g.

- 5) $(\forall \alpha < \omega_1)(\forall v \in d_\alpha \setminus d)(v \notin \bigcup_{w \in d} F(w))$.

Apply lemma 2.2 for the family $\{d_\alpha \setminus d : \alpha < \omega_1\}$ to get $\beta < \omega_1$ and an infinite set $A \subseteq \beta$ such that

- 6) $(\forall \alpha \in A)(\forall v \in d_\alpha \setminus d)(\forall w \in d_\beta \setminus d)(v \notin F(w))$.

Let $y \in p_\beta$. As in the claim the set

$$\{x \in \bigcup_{\alpha \in A} p_\alpha : \pi_2(x) \in d \text{ \& } x <_\Gamma y\}$$

is finite. Consequently we find $\alpha \in A$ such that

- 7) $(\forall x \in p_\alpha)(\forall y \in p_\beta)(\pi_2(x) \in d \Rightarrow \neg x <_\Gamma y)$.

We claim that p_α and p_β are compatible. Let $x \in p_\alpha$ and $y \in p_\beta$. By 7 we have that if $\pi_2(x) \in d$ then $\neg x <_\Gamma y$. If $\pi_2(x) \notin d$ and $\pi_2(y) \notin d$ then 6 applies and we get $\pi_2(x) \notin F(\pi_2(y))$. Finally if $\pi_2(x) \notin d$ and $\pi_2(y) \in d$ then we use 5 to conclude that $\pi_2(x) \notin F(\pi_2(y))$. Hence $x \in p_\alpha$, $y \in p_\beta$ and $x <_\Gamma y$ imply $\pi_2(x) \notin F(\pi_2(y))$. Consequently $p_\alpha \cup p_\beta \in \mathbf{P}_1$. ■

Lemma 2.6 *Assume that there exists a sequence $\{x_\alpha : \alpha < \omega_1\}$ of elements of ω^ω such that*

$$(\forall A \in [\omega_1]^{\omega_1})(\exists \alpha, \beta \in A)(\alpha < \beta \ \& \ x_\alpha \in F(x_\beta)).$$

Then the forcing notion $\mathbf{P}_1$ does not satisfy the Knaster condition.

PROOF For $\alpha < \omega_1$ choose $y_\alpha \in \Gamma$ such that

- $A_{y_\alpha} = \{\pi_0(y_\alpha)(n) : n \in \omega\} = \{x_\gamma : \gamma \leq \alpha\}$
- $\pi_1(y_\alpha)$ is the natural order on A_{y_α} , $x_\gamma <_{\pi_1(y_\alpha)} x_\beta$ iff $\gamma < \beta$.
- $\pi_2(y_\alpha) = x_\alpha$.

Let $p_\alpha = \{y_\alpha\}$ for $\alpha < \omega_1$. Then $\{p_\alpha : \alpha < \omega_1\}$ does not have an uncountable subset of pairwise compatible elements. ■

Putting together lemmas 2.5, 2.6 and 2.4 we get

Theorem 2.7 *It is consistent that there exists a ccc Souslin forcing notion which does not satisfy the Knaster condition.* ■

It is not difficult to see that this example does not satisfy the following requirement:

“The generic object is encoded by a real”

The next theorem says that also we can require such a condition. This answers a question of J. Bagaria.

Theorem 2.8 *It is consistent that there exists a ccc Souslin forcing notion $\mathbf{Q}$ such that $\Vdash_{\mathbf{Q}} \mathbf{V}[G] = \mathbf{V}[\dot{r}]$ for some $\mathbf{Q}$ -name $\dot{r}$ for a real and $\mathbf{Q}$ does not satisfy the Knaster condition.*

PROOF We follow the notation of the previous results. We work in the model of 2.7. Let $\mathbf{Q} = \{(p, w) : p \in \mathbf{P} \text{ \& } w \in [\omega^{<\omega}]^{<\omega}\}$ be ordered by

$$(p, w) \leq (q, v) \text{ if and only if } \\ p \leq q, w \subseteq v \text{ and } x|n \notin v \setminus w \text{ for every } x \in p \text{ and } n \in \omega.$$

$\mathbf{Q}$ may be easily represented as a Souslin forcing notion (remember that “for each x in p ” is a quantification on natural numbers). Note that if $p \leq q$, $p, q \in \mathbf{P}$ and $w \in [\omega^{<\omega}]^{<\omega}$ then $(p, w) \leq (q, w)$. Hence $\mathbf{Q}$ satisfies the countable chain condition. If $\mathbf{Q}$ satisfied the Knaster condition then $\mathbf{P}$ would have satisfied it. We show that the $\mathbf{Q}$ -generic object is encoded by a real. Let $\dot{r}$ be a $\mathbf{Q}$ -name for a subset of $\omega^{<\omega}$ (a real) such that for any $\mathbf{Q}$ -generic G we have $\dot{r}^G = \bigcup \{w : (\exists p)((p, w) \in G)\}$. Now in $\mathbf{V}[\dot{r}^G]$ define

$$H = \{(p, w) \in \mathbf{Q} : w \subseteq \dot{r}^G \text{ \& } (\forall x \in p)(\forall n \in \omega)(x|n \in \dot{r}^G \Leftrightarrow x|n \in w)\}.$$

Note that H includes G since $x \in p$, $x|n \notin w$ imply $(p, w) \Vdash x|n \notin \dot{r}$. H is a filter - suppose $(p_0, w_0), (p_1, w_1) \in H$. For each $x \in p_0 \cup p_1$ we find $(p_x, w_x) \in G$ such that $(p_x, w_x) \Vdash (\forall n \geq N)(x|n \notin \dot{r})$ for some N . If $x \notin p_x$ we could take large n and add $x|n$ to w_x . Then we would have $(p_x, w_x) \leq (p_x, w_x \cup \{x|n\})$ and $(p_x, w_x \cup \{x|n\}) \Vdash x|n \in \dot{r}$. Thus $x \in p_x$ for all $x \in p_0 \cup p_1$. Let $p = \bigcup_{x \in p_0 \cup p_1} p_x$, $w = \bigcup_{x \in p_0 \cup p_1} w_x$. Then $(p, w) \in G \subseteq H$ and $(p_0, w_0), (p_1, w_1) \leq (p, w)$. Consequently $H = G$ and the theorem is proved. ■

In the same time when the forcing notion $\mathbf{P}_1$ was constructed S.Todorćević found another example of this kind.

Let $\mathcal{F}$ be the family of all converging sequences s of real numbers such that $\lim s \notin s$. Todorćević’s forcing notion $\mathbf{P}_1^*$ consists of finite subsets p of $\mathcal{F}$ with property that

$$(\forall s, t \in p)(s \neq t \Rightarrow \lim s \notin t).$$

Todorćević proved that $\mathbf{P}_1^*$ satisfies ccc and that if $\mathfrak{b} = \omega_1$ then $\mathbf{P}_1^*$ does not have Knaster property (see [To]).

3 A nonhomogeneous example

In this section we give an example of ccc Souslin forcing notion which is very nonhomogeneous. Our forcing $\mathbf{P}_2$ will satisfy the following property:

there is no $p \in \mathbf{P}_2$ such that $p \Vdash \text{“}\hat{\mathbf{P}}_2|p \text{ is } \sigma\text{-centered”}$.

Recall that if $\mathbf{Q}$ is the Amoeba Algebra for Measure or the Measure Algebra then $\Vdash_{\mathbf{Q}} \text{“}\hat{\mathbf{Q}} \text{ is } \sigma\text{-centered”}$ (see [BJ]). The Todorćević example $\mathbf{P}_1^*$ has this property too.

Proposition 3.1 $\Vdash_{\mathbf{P}_1^*} \text{“}\hat{\mathbf{P}}_1^* \text{ is } \sigma\text{-centered”}$

PROOF For a rational number $d \in Q$ let $\phi_d : \mathbf{P}_1^* \longrightarrow \mathbf{P}_1^*$ be the translation by d . Thus $\phi_d(p) = \{s + d : s \in p\}$. Note that ϕ_d is an automorphism of $\mathbf{P}_1^*$. Moreover if $p_1, p_2 \in \mathbf{P}_1^*$ then $\bigcup \{s - r : s \in p_1, r \in p_2\}$ is a nowhere dense set. Hence we find a rational d such that

$$-d \notin \{a - b : a \in s \cup \{\lim s\}, b \in r \cup \{\lim r\}, s \in p_1, r \in p_2\}.$$

Then the conditions $\phi_d(p_1)$ and p_2 are compatible. Thus we have proved that for each $p \in \mathbf{P}_1^*$ the set $\{\phi_d(p) : d \in Q\}$ is predense in $\mathbf{P}_1^*$. This implies that

$$\Vdash_{\mathbf{P}_1^*} \text{“}\hat{\mathbf{P}}_1^* = \bigcup_{d \in Q} \phi_d[\dot{\Gamma}] \text{ and each } \phi_d[\dot{\Gamma}] \text{ is centered”},$$

where $\dot{\Gamma}$ is the canonical name for a generic filter. $\blacksquare$

We do not know if

$$\Vdash_{\mathbf{P}_1} \text{“}\hat{\mathbf{P}}_1 \text{ is } \sigma\text{-centered”}.$$

One can easily construct a ccc Souslin forcing $\mathbf{P}$ which does not force that $\hat{\mathbf{P}}$ is σ -centered. An example of such a forcing notion is the disjoint union of Cohen forcing and the measure algebra, $\mathbf{P} = (\{0\} \times \mathbf{C}) \cup (\{1\} \times \mathbf{B})$. In this order we have $(0, \emptyset) \Vdash \text{“}\{1\} \times \hat{\mathbf{B}} \text{ is not } \sigma\text{-centered”}$. But in this example we can find a dense set of conditions $p \in \mathbf{P}$ such that

$$p \Vdash_{\mathbf{P}} \text{“}\hat{\mathbf{P}}|p = \{q \in \hat{\mathbf{P}} : q \geq p\} \text{ is } \sigma\text{-centered”}.$$

Define $T^* \subseteq \omega^{<\omega}$, $f, g : T^* \longrightarrow \omega$ in such a way that:

- (α) T^* is a tree,
- (β) if $\eta \in T^*$ then $\text{succ}_{T^*}(\eta) = f(\eta)$,

- (γ) if $\text{lh}\eta < \text{lh}\nu$ or $\text{lh}\nu = \text{lh}\eta$ but $(\exists k < \text{lh}\eta)(\eta|k = \nu|k \ \& \ \eta(k) < \nu(k))$ then $f(\eta) < f(\nu)$,
- (δ) $g(\eta) > |T^* \cap \omega^{\text{lh}\eta}| \cdot \prod\{f(\nu) : f(\nu) < f(\eta)\} \cdot (100 + \text{lh}\eta)$,
- (ϵ) $f(\eta) > g(\eta) \cdot \prod\{2^{f(\nu)} : f(\nu) < f(\eta)\}$

For $\eta \in T^*$ and a set $A \subseteq \text{succ}_{T^*}(\eta) = f(\eta)$ we define a norm of A :

$$\mathbf{nor}_\eta(A) = \frac{g(\eta)}{|f(\eta) \setminus A|}.$$

Lemma 3.2 Suppose that $\eta \in T^*$ and $A_l \subseteq f(\eta)$ for $l < m$. Let $\zeta = \min\{\mathbf{nor}_\eta(A_l) : l < m\}$. Then

- 1) $\mathbf{nor}_\eta(\bigcap_{l < m} A_l) \geq \zeta/m$,
- 2) if $\zeta \geq 1$ and $m \leq \prod\{2^{f(\nu)} : f(\nu) < f(\eta)\}$ then $\bigcap_{l < m} A_l \neq \emptyset$.

PROOF 1) Note that

$$|f(\eta) \setminus \bigcap_{l < m} A_l| = \left| \bigcup_{l < m} f(\eta) \setminus A_l \right| \leq \sum_{l < m} |f(\eta) \setminus A_l| \leq m \cdot g(\eta)/\zeta$$

Hence

$$\mathbf{nor}_\eta\left(\bigcap_{l < m} A_l\right) = \frac{g(\eta)}{|f(\eta) \setminus \bigcap_{l < m} A_l|} \geq \zeta/m.$$

2) Applying 1) we get $\mathbf{nor}_\eta(\bigcap_{l < m} A_l) \geq 1/m$. Hence

$$|f(\eta) \setminus \bigcap_{l < m} A_l| \leq g(\eta) \cdot m \leq g(\eta) \cdot \prod\{2^{f(\nu)} : f(\nu) < f(\eta)\} < f(\eta)$$

(the last inequality is guaranteed by condition (ϵ)). Consequently the set $\bigcap_{l < m} A_l$ is nonempty. ■

Let $\mathbf{P}_2$ consists of all trees $T \subseteq T^*$ such that

$$\lim_{n \rightarrow \infty} \min\{\mathbf{nor}_\eta(\text{succ}_T(\eta)) : \eta \in T \cap \omega^n\} = \infty.$$

The order is the inclusion.

Recall that a forcing notion $\mathbf{Q}$ is σ - k -linked if there exist sets $R_n \subseteq \mathbf{Q}$ (for $n \in \omega$) such that $\bigcup_{n \in \omega} R_n = \mathbf{Q}$ and each R_n is k -linked (i.e. any k members of R_n has a common upper bound in $\mathbf{Q}$).

Proposition 3.3 *For every $k < \omega$ the forcing notion $\mathbf{P}_2$ is σ - k -linked.*

PROOF Let $n \in \omega$ be such that for each $\eta \in T^* \cap \omega^n$

$$k < \prod \{2^f(\nu) : f(\nu) < f(\eta)\}.$$

Note that the set

$$\{T \in \mathbf{P}_2 : \text{lh}(\text{root}T) \geq n \ \& \ (\forall \eta \in T)(\text{root}T \subseteq \eta \Rightarrow \mathbf{nor}_\eta(\text{succ}_T(\eta)) \geq 1)\}$$

is dense in $\mathbf{P}_2$. For $\eta \in T^*$, $\text{lh}\eta \geq n$ define

$$\mathcal{D}_\eta = \{T \in \mathbf{P}_2 : \text{root}T = \eta \ \& \ (\forall \nu \in T)(\eta \subseteq \nu \Rightarrow \mathbf{nor}_\nu(\text{succ}_T(\nu)) \geq 1)\}.$$

Since $\bigcup \{\mathcal{D}_\eta : \text{lh}\eta \geq n\}$ is dense in $\mathbf{P}_2$ it is enough to show that each $\mathcal{D}_\eta$ is k -linked. Suppose $T_0, \dots, T_{k-1} \in \mathcal{D}_\eta$. Since $k < \prod \{2^f(\nu) : f(\nu) < f(\eta)\}$ we may apply lemma 3.2 2) to conclude that if $\nu \in T = T_0 \cap \dots \cap T_{k-1}$, $\nu \supseteq \eta$ then $\text{succ}_T(\nu) \neq \emptyset$. By 3.2 1) we get $T \in \mathbf{P}_2$. ■

For $\eta \in T^*$ we define the forcing notion $\mathbf{Q}_\eta$:

$$\begin{aligned} \mathbf{Q}_\eta = \{t \subseteq T^* : & \ t \text{ is a finite tree of the height } n \in \omega, \\ & \text{root}t = \eta \text{ and} \\ & (\forall \nu \in t \cap \omega^{< n})(\eta \subseteq \nu \Rightarrow \mathbf{nor}_\nu(\text{succ}_t(\nu)) \geq \text{lh}\nu)\}. \end{aligned}$$

Since $\mathbf{Q}_\eta$ is countable and atomless it is isomorphic to Cohen forcing $\mathbf{C}$. Let $\mathbf{P} = \prod \{\mathbf{Q}_{i,\eta} : i < \omega_1, \eta \in T^*\}$ be the finite support product such that each $\mathbf{Q}_{i,\eta}$ is a copy of $\mathbf{Q}_\eta$.

Theorem 3.4 *Let $G \subseteq \mathbf{P}$ be a generic filter over $\mathbf{V}$. Then, in $\mathbf{V}[G]$, there is no $S \in \mathbf{P}_2$ such that*

$$S \Vdash_{\mathbf{P}_2} \text{“}\hat{\mathbf{P}}_2|S \text{ is } \sigma\text{-centered”}.$$

PROOF We work in $\mathbf{V}[G]$.

Assume $S \Vdash \text{“}\hat{\mathbf{P}}_2|S \text{ is } \sigma\text{-centered”}$. Let $\dot{\mathcal{R}}_n$ ($n \in \omega$) be $\mathbf{P}_2$ -names for subsets of $\mathbf{P}_2$ such that

$$S \Vdash_{\mathbf{P}_2} \text{“}\hat{\mathbf{P}}_2|S \subseteq \bigcup_{n \in \omega} \dot{\mathcal{R}}_n \ \& \ \text{each } \dot{\mathcal{R}}_n \text{ is directed”}.$$

Take $n \in \omega$ such that

$$(\forall \eta \in S)(n \leq \text{lh} \eta \Rightarrow \mathbf{nor}_\eta(\text{succ}_S(\eta)) \geq 1).$$

Fix any $\eta \in S \cap \omega^n$ and choose $l, m \in \text{succ}_S(\eta)$, $l < m$. For $i < \omega_1$ put

$$T_i = \bigcup \{t : \{(i, \eta \hat{m}), t\} \in G\}.$$

Each T_i is the tree added by $G \cap \mathbf{Q}_{i, \eta \hat{m}}$ and it is an element of $\mathbf{P}_2$. Moreover $\text{root} T_i = \eta \hat{m}$ and for each $\nu \in T_i$ if $\eta \hat{m} \subseteq \nu$ then $\mathbf{nor}_\nu(\text{succ}_{T_i}(\nu)) \geq \text{lh} \nu$. Hence, by lemma 3.2, $T_i \cap S \in \mathbf{P}_2$ for each $i \in \omega_1$.

Now we work in $\mathbf{V}$.

We find $p_i, \dot{S}_i, \eta_i, n_i$ such that for each $i \in \omega_1$:

- $p_i \in \mathbf{P}, n_i \in \omega, \eta_i \in T^*$ and $\dot{S}_i$ is a $\mathbf{P}$ -name for a member of $\mathbf{P}_2$,
- $\Vdash_{\mathbf{P}} (\forall \nu \in \dot{S}_i)(\text{root} \dot{S}_i \subseteq \nu \Rightarrow \mathbf{nor}_\nu(\text{succ}_{\dot{S}_i}(\nu)) \geq 1)$,
- $p_i \Vdash_{\mathbf{P}} "\eta \hat{l} \subseteq \eta_i = \text{root} \dot{S}_i \ \& \ \dot{S}_i \Vdash_{\mathbf{P}_2} \dot{T}_i \in \dot{R}_{n_i}"$,
- $(i, \eta \hat{m}) \in \text{dom} p_i$.

Next we find a set $I \in [\omega_1]^{\omega_1}$ such that $\{\text{dom} p_i : i \in I\}$ forms a Δ -system with the root d and for each $i \in I$:

- $\eta_i = \eta^*$ and $n_i = n^*$,
- $p_i|d = p^*$,
- $p_i(i, \eta \hat{m}) = t, (i, \eta \hat{m}) \notin d$.

Let $n^\#$ be the height of the tree t . Clearly we may assume that $n^\# > \text{lh} \eta^*$.

Fix an enumeration $\{\rho_k : k < k^\#\}$ of $t \cap \omega^{n^\#}$. Put

$$H = \{(a_k : k < k^\#) : a_k \subseteq f(\rho_k) \ \& \ \mathbf{nor}_{\rho_k}(a_k) \geq n^\#\}.$$

Choose distinct $i_{\bar{a}} \in I$ for $\bar{a} \in H$. We define a condition $q \in \mathbf{P}$ extending all $p_{i_{\bar{a}}}$ ($\bar{a} \in H$):

$$\begin{aligned} \text{dom} q &= \bigcup \{\text{dom} p_{i_{\bar{a}}} : \bar{a} \in H\}; \\ \text{if } (i, \nu) \in \text{dom} p_{i_{\bar{a}}}, (i, \nu) \neq (i_{\bar{a}}, \eta \hat{m}) \text{ then } q(i, \nu) &= p_{i_{\bar{a}}}(i, \nu); \\ q(i_{\bar{a}}, \eta \hat{m}) &= t \cup \{\rho_k \hat{c} : k < k^\#, c \in \bar{a}(k)\}. \end{aligned}$$

Now we take $r \geq q$ such that r decides all $\dot{S}_{i_{\bar{a}}}|(n^\# + 1)$. Thus we have finite trees $s_{\bar{a}}$ (for $\bar{a} \in H$) such that $r \Vdash_{\mathbf{P}} \dot{S}_{i_{\bar{a}}}|(n^\# + 1) = s_{\bar{a}}$.

CLAIM: : There exists $H' \subseteq H$ such that

- (i) $\bigcap_{\bar{a} \in H'} (s_{\bar{a}} \cap \omega^{n^\#} + 1) \neq \emptyset$ and
- (ii) for each $k < k^\#$ the set $\bigcap \{\bar{a}(k) : \bar{a} \in H'\}$ is empty.

Indeed, let $H_\rho = \{\bar{a} \in H : \rho \in s_{\bar{a}}\}$ for $\rho \in T^* \cap \omega^{n^\#} + 1$, $\eta \hat{l} \subseteq \rho$. Clearly $\rho \in \bigcap_{\bar{a} \in H_\rho} s_{\bar{a}}$, so it is enough to show that for some ρ the family H_ρ satisfies (ii). Suppose that for each $\rho \in T^* \cap \omega^{n^\#} + 1$, $\rho \supseteq \eta \hat{l}$ we can find $k_\rho < k^\#$ and c_ρ such that $c_\rho \in \bigcap \{\bar{a}(k_\rho) : \bar{a} \in H_\rho\}$. Put

$$\bar{a}^*(k) = f(\rho_k) \setminus \{c_\rho : \rho \in T^* \cap \omega^{n^\#} + 1 \text{ \& } \eta \hat{l} \subseteq \rho\}.$$

Let $\rho^+ \in T^* \cap \omega^{n^\#}$ be such that

$$f(\rho^+) = \max\{f(\rho) : \rho \in T^* \cap \omega^{n^\#}, \eta \hat{l} \subseteq \rho\}.$$

By condition (γ) we get

$$|\{\rho \in T^* \cap \omega^{n^\#} + 1 : \eta \hat{l} \subseteq \rho\}| \leq \prod \{f(\nu) : f(\nu) \leq f(\rho^+)\}.$$

Now, for each $k < k^\#$ we have $f(\rho^+) < f(\rho_k)$ (recall that $\eta \hat{l} \subseteq \rho^+$, $\eta \hat{m} \subseteq \rho_k$ and $l < m$ so condition (γ) works). Hence

$$\mathbf{nor}_{\rho_k}(\bar{a}^*(k)) \geq \frac{g(\rho_k)}{\prod \{f(\nu) : f(\nu) \leq f(\rho^+)\}} \geq \frac{g(\rho_k)}{\prod \{f(\nu) : f(\nu) < f(\rho_k)\}} > n^\#$$

Thus $\bar{a}^* \in H$. Since $c_\rho \notin \bar{a}^*(k_\rho)$ we have $\bar{a}^* \notin H_\rho$ for every $\rho \in T^* \cap \omega^{n^\#} + 1$, $\eta \hat{l} \subseteq \rho$. Since $\bigcup \{H_\rho : \rho \in T^* \cap \omega^{n^\#} + 1, \eta \hat{l} \subseteq \rho\} = H$ we get a contradiction. The claim is proved.

Now let $H' \subseteq H$ be a family given by the claim. Condition (ii) implies that

$$r \Vdash_{\mathbf{P}} \text{“the family } \{T_{i_{\bar{a}}} : \bar{a} \in H'\} \text{ has no upper bound in } \mathbf{P}_2\text{”}.$$

Since $|H| \leq \prod \{2^{f(\rho_k)} : k < k^\#\}$ we have that for each $\rho \in T^* \cap \omega^{n^\#} + 1$

$$|H'| \leq \prod \{2^{f(\nu)} : f(\nu) < f(\rho)\}.$$

Hence we may apply 3.2 2) to conclude that for every $\rho \supseteq \eta^l$, $\text{lh}\rho \geq n^\# + 1$:

$$r \Vdash_{\mathbf{P}} \text{“if } \rho \in \bigcap_{\bar{\mathbf{a}} \in H'} \dot{S}_{i_{\bar{\mathbf{a}}}} \text{ then } \bigcap_{\bar{\mathbf{a}} \in H'} \text{succ}_{\dot{S}_{i_{\bar{\mathbf{a}}}}}(\rho) \neq \emptyset\text{”}.$$

Thus

$$r \Vdash_{\mathbf{P}} \text{“the family } \{\dot{S}_{i_{\bar{\mathbf{a}}}} : \bar{\mathbf{a}} \in H'\} \text{ has an upper bound”}.$$

Since $r \Vdash_{\mathbf{P}} \text{“}\dot{S}_{i_{\bar{\mathbf{a}}}} \Vdash_{\mathbf{P}_2} \dot{T}_{i_{\bar{\mathbf{a}}}} \in \dot{R}_{n^*}\text{”}$ we get a contradiction. ■

REMARK: 1) In the above theorem we worked in the model $\mathbf{V}[G]$ for technical reasons only. The assertion of the theorem can be proved in ZFC.

2) The forcing notion $\mathbf{P}_2$ is a special case of the forcing studied in [Sh1].

Problem 3.5 *Does there exist a ccc Souslin forcing $\mathbf{P}$ such that*

- 1) $\mathbf{P}$ is homogeneous (i.e. for each $p \in \mathbf{P}$, $\Vdash_{\mathbf{P}}$ “there exists a generic filter G over $\mathbf{V}$ such that $p \in G$ ”)
- 2) $\nVdash_{\mathbf{P}} \text{“}\dot{\mathbf{P}} \text{ is } \sigma\text{-centered”}$?

4 On “small subsets of $\mathbf{P}$ are σ -centered”.

Our next example is connected with the following, still open, question:

Problem 4.1 *Assume that for each ccc Souslin forcing $\mathbf{P}$ every set $Q \in [\mathbf{P}]^{\omega_1}$ is σ -centered (in $\mathbf{P}$).*

Does $\mathbf{MA}_{\omega_1}(\text{Souslin})$ hold true?

As an illustration of this subject let us recall a property of Random (Solovay) Algebra $\mathbf{B}$ (see [BaJ]):

if every $B \in [\mathbf{B}]^{\omega_1}$ is σ -centered
then the real line can not be covered by ω_1 null sets and consequently $\mathbf{MA}_{\omega_1}(\mathbf{B})$ holds true.

Our example shows that the above property of the algebra $\mathbf{B}$ does not extend for other forcing notions. Let

$$\mathbf{P}_3 = \{(n, T) : n \in \omega \ \& \ T \subseteq 2^{<\omega} \text{ is a tree} \ \& \ (\forall t \in T \cap 2^n)(\mu([T_t]) > 0)\}.$$

The order is defined by

$(n_1, T_1) \leq (n_2, T_2)$ if and only if
 $n_1 \leq n_2$, $T_2 \subseteq T_1$ and $T_1|_{n_1} = T_2|_{n_1}$.

Lemma 4.2 $\mathbf{P}_3$ is a σ -linked Souslin forcing which is not σ -centered.

PROOF Note that $\Vdash_{\mathbf{P}_3}$ “there exists a perfect set of random reals over $\mathbf{V}$ ”. Hence $\mathbf{P}_3$ is not σ -centered. To show that it is σ -linked define sets $U(W, n, m)$ for $n < m < \omega$ and finite trees $W \subseteq 2^{\leq m}$:

$$U(W, n, m) = \{(n, T) \in \mathbf{P}_3 : T|_m = W \text{ \& } (\forall t \in T \cap 2^n)(\mu([T_t]) > W(t)/2^{m+1})\}$$

where $W(t) = |\{s \in W \cap 2^m : t \subseteq s\}|$ (for $t \in W \cap 2^n$). Clearly each set $U(W, n, m)$ is linked (i.e. each two members of it are compatible in $\mathbf{P}_3$) and $\mathbf{P}_3 = \bigcup \{U(W, n, m) : n < m < \omega \text{ \& } W \subseteq 2^{\leq m}\}$. Since obviously $\mathbf{P}_3$ is Souslin we are done. ■

Let $\mathbf{B}(\kappa)$ stand for Random Algebra for adding κ many random reals. This is the measure algebra of the space 2^κ .

Theorem 4.3 Assume $\mathbf{V} \models \mathbf{CH}$. Let $G \subseteq \mathbf{B}(\omega_2)$ be a generic set over $\mathbf{V}$. Then, in $\mathbf{V}[G]$

- (i) Martin axiom fails for $\mathbf{P}_3$ but
- (ii) each $Q \in [\mathbf{P}_3]^{\omega_1}$ is σ -centered (in $\mathbf{P}_3$).

PROOF Cichon proved that one random real does not produce a perfect set of random reals (see [BaJ]). Hence in $\mathbf{V}[G]$ there is no perfect set of random reals over $\mathbf{V}$. Consequently the first assertion is satisfied in $\mathbf{V}[G]$. Since $\mathbf{V}[G] \models$ “each $B \in [\mathbf{B}]^{\omega_1}$ is σ -centered in $\mathbf{B}$ ” (compare section 3) it is enough to show the following

CLAIM: Suppose that each $B \in [\mathbf{B}]^{\omega_1}$ is σ -centered. Then every set $Q \in [\mathbf{P}_3]^{\omega_1}$ is σ -centered.

Indeed, let $Q \in [\mathbf{P}_3]^{\omega_1}$. For $n \in \omega$ and $t \in 2^n$ put

$$B(t, n) = \{[T_t] : (n, T) \in Q \text{ \& } t \in T\}.$$

By our assumption we find sets $B(t, n, k)$ for $k, n \in \omega$, $t \in 2^n$ such that $B(t, n) = \bigcup_{k \in \omega} B(t, n, k)$ and for each $A_1, A_2 \in B(t, n, k)$ the set $A_1 \cap A_2$

is of positive measure. Now define sets $Q(n, W, \sigma)$ for $n \in \omega$, a finite tree $W \subseteq 2^{\leq n}$ and a function $\sigma : W \cap 2^n \rightarrow \omega$:

$$Q(n, W, \sigma) = \{(n, T) \in Q : T|n = W \text{ \& } (\forall t \in T \cap 2^n)([T_t] \in B(t, n, \sigma(t)))\}.$$

Note that if $(n, T_1), (n, T_2) \in Q(n, W, \sigma)$ then for each $t \in W \cap 2^n$ the set $[(T_1)_t] \cap [(T_2)_t]$ is of positive measure. Consequently each $Q(n, W, \sigma)$ is linked and we are done. ■

5 A σ -centered example

In this section we define a very simple σ -centered Souslin forcing notion. Next we show that in any generic extension of some model of **CH** via finite support iteration of the Dominating (Hechler) Algebra, Martin Axiom fails for this forcing notion. Consequently we get the consistency of the following sentence:

*any union of less than continuum meager sets is meager + $\neg$ CH
+ **MA** fails for some σ -centered Souslin forcing.*

Our example $\mathbf{P}_4$ consists of all pairs (n, F) such that $n \in \omega$, $F \in [2^\omega]^{<\omega}$ and all elements of the list $\{x|n : x \in F\}$ are distinct. $\mathbf{P}_4$ is ordered by

$$(n, F) \leq (n', F') \text{ if and only if} \\ n \leq n', F \subseteq F' \text{ and } \{x|n : x \in F\} = \{x|n : x \in F'\}.$$

Lemma 5.1 $\mathbf{P}_4$ is a σ -centered Souslin forcing.

PROOF Clearly $\mathbf{P}_4$ is Souslin (even Borel). To show that $\mathbf{P}_4$ is σ -centered note that if $\{x|n : x \in F_0\} = \dots = \{x|n : x \in F_k\}$ then the conditions $(n, F_0), \dots, (n, F_k)$ are compatible (if m is large enough then $(m, F_0 \cup \dots \cup F_k)$ is a witness for this). ■

Now we want to define the model we will start with. At the beginnig we work in **L**. Applying the technology of [Sh] we can construct a sequence $(\mathbf{P}_\xi : \xi \leq \omega_1)$ of forcing notions such that for each $\alpha, \beta < \omega_1$, $\xi \leq \omega_1$:

(1) if $\alpha < \beta$ then $\mathbf{P}_\alpha$ is a complete suborder of $\mathbf{P}_\beta$,

- (2) there is $\gamma > \beta$ such that $\mathbf{P}_{\gamma+1} = \mathbf{P}_\gamma * \dot{\mathbf{D}}_\alpha$, where $\dot{\mathbf{D}}_\alpha$ is the $\mathbf{P}_\gamma$ -name for finite support, α in length, iteration of Hechler forcing,
- (3) $\mathbf{P}_\xi$ satisfies ccc,
- (4) if ξ is limit then $\mathbf{P}_\xi = \overrightarrow{\lim_{\zeta < \xi}} \mathbf{P}_\zeta$,
- (5) $\mathbf{P}_{\omega_1} \Vdash$ “every projective set of reals has Baire property”

(for details see also [JR]). Recall that Hechler forcing $\mathbf{D}$ consists of all pairs (n, f) such that $n \in \omega$, $f \in \omega^\omega$. These pairs are ordered by

$$(n, f) \leq (n', f') \text{ if and only if } n \leq n', f|n = f'|n' \text{ and } f(k) \leq f'(k) \text{ for all } k \in \omega.$$

Suppose $G \subseteq \mathbf{P}_{\omega_1}$ is a generic set over $\mathbf{L}$. We work in $\mathbf{L}[G]$. For distinct $x, y \in 2^\omega$ we define $h(x, y) = \min\{n : x(n) \neq y(n)\}$. Easy calculations show the following

Lemma 5.2 *Let $b \subseteq \omega$. Then the following conditions are equivalent:*

- (i) *there exists a Borel equivalence relation R on 2^ω with countable many equivalence classes such that $\{h(x, y) : x, y \in 2^\omega \cap \mathbf{L} \text{ \& } x \neq y \text{ \& } R(x, y)\} \subseteq b$,*
- (ii) *there exists an equivalence relation R on 2^ω with countable many equivalence classes such that $\{h(x, y) : x, y \in 2^\omega \cap \mathbf{L} \text{ \& } x \neq y \text{ \& } R(x, y)\} \subseteq b$,*
- (iii) *there exist sets $Y_n \subseteq 2^\omega$ (for $n \in \omega$) such that $\mathbf{L} \cap 2^\omega \subseteq \bigcup_{n \in \omega} Y_n$ and $\bigcup_{n \in \omega} \{h(x, y) : x \neq y \text{ \& } x, y \in Y_n\} \subseteq b$,*
- (iv) $(\exists f : 2^{<\omega} \rightarrow 2)(\forall x \in 2^\omega \cap \mathbf{L})(\exists m \in \omega)(\forall n > m)(n \notin b \Rightarrow f(x|n) = x(n)). \blacksquare$

The Raisonniier filter $\mathcal{F}$ consists of all sets $b \subseteq \omega$ satisfying one of the conditions of 5.2 (cf [Ra]). $\mathcal{F}$ is a proper filter on ω . Directly from (iv) of 5.2 one can see that $\mathcal{F}$ is a Σ_3^1 -subset of 2^ω . Consequently it has Baire property (recall that we are in $\mathbf{L}[G]$).

Theorem 5.3 (Talagrand, [Ta]) *For any proper filter F on ω the following conditions are equivalent:*

- (i) *F does not have Baire property,*

(ii) for every increasing sequence $(n_k : k \in \omega)$ of integers there exists $b \in F$ such that $(\exists_k^\infty)(b \cap [n_k, n_{k+1}) = \emptyset)$. ■

Applying the above theorem we can find an increasing function $r \in \omega^\omega \cap \mathbf{L}[G]$ such that (in $\mathbf{L}[G]$)

$$(\forall b \in \mathcal{F})(\forall_k^\infty)(b \cap [r(k), r(k+1)) \neq \emptyset)$$

Let $\dot{r}$ be the $\mathbf{P}_{\omega_1}$ -name for r and let $\alpha_0 < \omega_1$ be such that $\dot{r}$ is a $\mathbf{P}_{\alpha_0}$ -name.

Our basic model will be $\mathbf{L}[\dot{r}]$

Theorem 5.4 *Let κ be a regular cardinal. Let $\mathbf{D}_\kappa$ be the finite support iteration of Hechler forcing of the length κ . Suppose $H \subseteq \mathbf{D}_\kappa$ is a generic set over $\mathbf{L}[\dot{r}]$. Then*

$$\mathbf{L}[\dot{r}][H] \models \text{“there is no } \mathbf{P}_4\text{-generic over } \mathbf{L}[\dot{r}]\text{.”}$$

PROOF Assume not. Let $H^* \in \mathbf{L}[\dot{r}][H]$ be a $\mathbf{P}_4$ -generic over $\mathbf{L}[\dot{r}]$. Put $T = \bigcup \{F : (\exists n \in \omega)((n, F) \in H^*)\}$. Then in $\mathbf{L}[\dot{r}][H^*]$ we have:

(6) T is a closed subset of 2^ω ,

(7) $(\exists_k^\infty)(\forall x, y \in T)(x \neq y \Rightarrow h(x, y) \notin [r(k), r(k+1)))$ and

(8) $(\forall x \in 2^\omega \cap \mathbf{L})(\exists q \in Q)(q + x \in T)$

(Q stands for the set of all sequences eventually equal 0, $+$ denotes the addition modulo 2). Since both (7) and (8) are absolute (Π_2^1) sentences they are satisfied in $\mathbf{L}[\dot{r}][H]$ too. Let $\dot{T} \in \mathbf{L}[\dot{r}]$ be a $\mathbf{P}_\kappa$ -name for T . Since T is a closed subset of 2^ω we can think of $\dot{T}$ as a name for a real.

Now we work in $\mathbf{L}[\dot{r}]$. Let $p \in \mathbf{D}_\kappa \cap \mathbf{L}[\dot{r}]$ be such that

$$p \Vdash \text{“}\dot{T} \text{ satisfies (6), (7) and (8)”}.$$

By Souslin forcing properties (see §1 of [JS1]) we find a (closed) countable set $S \subseteq \kappa$ such that:

(9) $\dot{T}$ is a $\mathbf{D}_\kappa|S$ -name, $p \in \mathbf{D}_\kappa|S$ and

(10) $\mathbf{D}_\kappa|S$ is a complete suborder of $\mathbf{D}_\kappa$.

Since (6)-(8) are absolute we get

$$(11) \quad p \Vdash_{\mathbf{D}_\kappa|S} \dot{T} \text{ satisfies (6), (7) and (8)}.$$

But $\mathbf{D}_\kappa|S$ is isomorphic to finite support iteration of Hechler forcing of the countable length α ($\alpha < \omega_1$). Thus we can treat $\dot{T}$ as a $\mathbf{D}_\alpha$ -name and p as a condition in $\mathbf{D}_\alpha$. Then, in $\mathbf{L}[r]$

$$(12) \quad p \Vdash_{\mathbf{D}_\alpha|S} \dot{T} \text{ satisfies (6), (7) and (8)}.$$

By (2) we find $\beta > \alpha_0$ such that

$$(13) \quad \mathbf{P}_{\gamma+1} = \mathbf{P}_\gamma * \dot{\mathbf{D}}_\alpha \text{ and}$$

$$(14) \quad p \equiv (1, p) \text{ interpreted as a member of } \mathbf{P}_{\gamma+1} \text{ belongs to } G.$$

By Souslin forcing properties (12) holds true in $\mathbf{L}[G \cap \mathbf{P}_\gamma]$ and hence

$$(15) \quad \mathbf{L}[G \cap \mathbf{P}_{\gamma+1}] \models \dot{T}^G \text{ satisfies (6), (7) and (8)}$$

(we treat here $\dot{T}$ as a $\mathbf{P}_{\gamma+1}$ -name). Let $b = \{h(x, y) : x \neq y \ \& \ x, y \in \dot{T}^G\} \in [\omega]^{<\omega} \cap \mathbf{L}[G]$. By (15) and by Shoenfield absoluteness we have

$$(16) \quad \mathbf{L}[G] \models \dot{T}^G \text{ satisfies (6), (7) and (8)}.$$

Since $\{h(x, y) : x \neq y \ \& \ x, y \in \dot{T}^G\} = \{h(x, y) : x \neq y \ \& \ x, y \in \dot{T}^G + q\}$ we conclude that

$$(17) \quad \mathbf{L}[G] \models \text{“sets } \dot{T}^G + q \text{ (for } q \in Q) \text{ witness that } b \in \mathcal{F} \text{” and}$$

$$(18) \quad \mathbf{L}[G] \models (\exists_k^\infty)(b \cap [r(k), r(k+1)) = \emptyset).$$

The last condition contradicts our choice of r . ■

Since $\Vdash_{\mathbf{D}_\kappa}$ “any union of less than κ meager sets is meager” we get

Corollary 5.5 *The following theory is consistent:*

ZFC + $\neg\mathbf{CH}$ + “Martin Axiom fails for some σ -centered Souslin forcing”
+ “any union of less than continuum meager sets is meager”. ■

6 On Souslin not ccc

In this section we will give a negative answer to the following question of Woodin:

If $\mathbf{P}$ is a Souslin forcing notion which is not ccc
then there exists a perfect set $T \subseteq \mathbf{P}$ such that each distinct
 $t_1, t_2 \in T$ are incompatible.

Recall that in the case of non-ccc partial orders we do not require Souslin forcings to satisfy the condition:

“the set $\{(p, q) : p \text{ is incompatible with } q\}$ is Σ_1^1 ”.

Thus a forcing notion $\mathbf{P}$ is *is Souslin not ccc* if both $\mathbf{P}$ and $\leq_{\mathbf{P}}$ are analytic sets. The reason for this is that we want to cover in our definition various standard forcing notions with simple definitions for which incompatibility is not analytic (e.g. Laver forcing).

Let $\mathbf{Q}$ be the following partially ordered set:

$W \in \mathbf{Q}$ if W is a finite set of pairs (α, β) , $\alpha \leq \beta < \omega_1$ such that if $(\alpha_1, \beta_1), (\alpha_2, \beta_2)$ are in W , then $\beta_1 < \alpha_2$ or $\beta_2 < \alpha_1$.

$\mathbf{Q}$ is ordered by the inclusion.

It follows from [Je1] that $\mathbf{Q}$ is proper. Clearly $|\mathbf{Q}| = \omega_1$. Next define a forcing notion $\mathbf{P}_5$. It consists of all $r \in \omega^\omega$ such that r codes a pair (E^r, w^r) where

1. E^r is a relation on ω such that $(\omega, E^r) \models \mathbf{ZFC}^-$ and E^r encodes all elements of $\omega \cup \{\omega\}$.
2. $w^r \in \omega$ and $E^r \models “w^r \in \mathbf{Q}”$.

We say that a one-to-one function $f \in \omega^\omega$ *interprets* E^{r_1} in E^{r_2} if there exists $n \in \omega$ such that $\text{rng}(f) = \{k \in \omega : E^{r_2}(k, n)\}$ and $E^{r_1}(l, k) \equiv E^{r_2}(f(l), f(k))$.

If f interprets E^{r_1} in E^{r_2} then E^{r_2} may “discover” that some of the ordinals of E^{r_1} are not ordinals (i.e. not well-founded). Let $w(r_1, r_2, f) = w^{r_1} \cap \{(\alpha, \beta) : \alpha \leq \beta \text{ are ordinals in } E^{r_2}\}$. Then, in E^{r_2} , $w(r_1, r_2, f)$ is an initial segment of w^{r_1} and it is in $\mathbf{Q}^{E^{r_2}}$.

Now we can define the order $\leq$ on $\mathbf{P}_5$:

$r_1 \leq r_2$ if and only if $r_1 = r_2$ or
there exists $f \in \omega^\omega$ which interprets E^{r_1} in E^{r_2} and such that

$$(\omega, E^{r_2}) \models w(r_1, r_2, f) \subseteq w^{r_2}.$$

Obviously both $\mathbf{P}_5$ and the order $\leq$ are Σ_1^1 -sets.

For $r \in \mathbf{P}_5$ we define $W(r)$ as $w^r \cap \{(\alpha, \beta) : \alpha \leq \beta \text{ are well founded}\}$. Note that $W(r_1) = W(r_2)$ implies r_1 and r_2 are equivalent in $\mathbf{P}_5$ (i.e they have the same compatible elements of $\mathbf{P}_5$). Consequently $\mathbf{Q}$ may be densely embedded into the complete Boolean algebra determined by $\mathbf{P}_5$. It follows from [Je1] that $\mathbf{P}$ is proper, it is Souslin and it does not satisfy the countable chain condition. Moreover, if $\omega_1 < 2^\omega$ then $\mathbf{P}$ does not contain a perfect set of pairwise incompatible elements (recall $|\mathbf{Q}| = \omega_1$). ■

An interesting question appears here:

Suppose $\mathbf{P}$ is ω -proper and Souslin.

Does there exists a perfect set of pairwise incompatible elements of $\mathbf{P}$?

The negative answer to this question is given by the following result.

Theorem 6.1 *Assume $\omega_1 < cf(2^\omega)$. There exists an ω -proper Souslin not ccc forcing notion $\mathbf{P}_5^*$ with no perfect set of pairwise incompatible elements.*

PROOF Let $\delta \leq \omega_1$ be additively indecomposable. Let $\mathbf{Q}^*$ be the order defined by:

$W \in \mathbf{Q}^*$ if and only if

W is a countable set of pairs (α, β) , $\alpha \leq \beta < \omega_1$ such that

- $(\alpha_1, \beta_1), (\alpha_2, \beta_2) \in W \Rightarrow \beta_1 < \alpha_2$ or $\beta_2 < \alpha_1$,
- $\{(\alpha, \beta) \in W : \alpha \neq \beta\}$ is finite,
- the order type of the set $\{\alpha : (\exists \beta)((\alpha, \beta) \in W)\}$ is less than δ .

$\mathbf{Q}^*$ is ordered by the inclusion.

It follows from Chapter XVII, §3 of [Sh 2] that $\mathbf{Q}^*$ is α -proper for each $\alpha < \omega_1$.

Now we can repeat the coding procedure that we applied to define the forcing notion $\mathbf{P}_5$. Thus we get the Souslin forcing notion $\mathbf{P}_5^*$ such that $\mathbf{Q}^*$ can be densely embedded in the Boolean algebra determined by $\mathbf{P}_5^*$.

For $W \in \mathbf{Q}^*$ let $\text{heart}(W) = \{(\alpha, \beta) \in W : \alpha \neq \beta\}$.

Assume that $\{(E^{r_\eta}, w^{r_\eta}) : \eta \in 2^\omega\} \subseteq \mathbf{P}_5^*$ is a perfect set of pairwise incompatible elements. Let W_η be the well-founded part of w^{r_η} . Since w.l.o.g. we can assume that $\sup\{\beta : (\exists \alpha)((\alpha, \beta) \in W_\eta)\}$ is constant and $\text{heart}(W_\eta)$ is constant we easily get a contradiction. ■

7 On ccc Σ_2^1

Souslin ccc notions of forcing are indestructible ccc (see [JS1]):

Suppose $\mathbf{P}$ is a ccc Souslin notion of forcing. Let $\mathbf{Q}$ be a ccc forcing notion. Then $\Vdash_{\mathbf{Q}} \hat{\mathbf{P}}$ is ccc ”.

The above property does not hold true for more complicated forcing notions. In this section we show that there may exist two ccc Σ_2^1 -notions of forcing $\mathbf{P}_6$ and $\mathbf{P}_6^*$ such that $\mathbf{P}_6 \times \mathbf{P}_6^*$ does not satisfy ccc.

We start with $\mathbf{V} = \mathbf{L}$. Let $\mathbf{Q}$ be a ccc notion of forcing such that

$$\Vdash_{\mathbf{Q}} \mathbf{MA} + \neg \mathbf{CH}$$

Let $G \subseteq \mathbf{Q}$ be a generic set over $\mathbf{L}$ and let r be a random real over $\mathbf{L}[G]$. Recall that by theorem of Roitman (cf [Ro]) we have $\mathbf{L}[G][r] \models \mathbf{MA}(\sigma\text{-centered})$.

Fix a sequence $(f_\alpha : \alpha < \omega_1) \in \mathbf{L}$ of one-to-one functions $f_\alpha : \alpha \xrightarrow{1-1} \omega$ and define in $\mathbf{L}[r]$ sets E_1, E_2 by

$$E_i = \{\{\alpha, \beta\} \in [\omega_1]^2 : \beta < \alpha \ \& \ r(f_\alpha(\beta)) = i\} \quad \text{for } i = 0, 1.$$

We define forcing notions $\mathbf{P}_6, \mathbf{P}_6^*$:

$$\mathbf{P}_6 = \{H \in [\omega_1]^2 : [H]^2 \subseteq E_0\},$$

$$\mathbf{P}_6^* = \{H \in [\omega_1]^2 : [H]^2 \subseteq E_1\}.$$

Orders are inclusions.

Both $\mathbf{P}_6$ and $\mathbf{P}_6^*$ are elements of $\mathbf{L}[r]$. Moreover they can be thought of as subsets of $\mathbf{L}[r] \cap 2^\omega$. Applying $\mathbf{MA}(\sigma\text{-centered})$ we get that (cf [Je]):

$$\mathbf{L}[G][r] \models \text{“any subset of } \mathbf{L}[r] \cap 2^\omega \text{ is a relative } \Sigma_2^0\text{-set”}.$$

Consequently

$$\mathbf{L}[G][r] \models \text{“any subset of } \mathbf{L}[r] \cap 2^\omega \text{ is } \Sigma_2^1\text{”}.$$

Thus $\mathbf{P}_6$ and $\mathbf{P}_6^*$ are Σ_2^1 -notions of forcing in $\mathbf{L}[G][r]$ (i.e. both $\mathbf{P}_6$, $\mathbf{P}_6^*$ and orders and the relations of incompatibility are Σ_2^1 -sets). Roitman proved the following

Theorem 7.1 (Roitman, Prop.4.6 of [Ro]) *In $\mathbf{L}[G][r]$ both $\mathbf{P}_6$ and $\mathbf{P}_6^*$ satisfy ccc and $\mathbf{P}_6 \times \mathbf{P}_6^*$ does not satisfy ccc.* ■

Corollary 7.2 *The following theory is consistent:*

$\mathbf{ZFC} + \mathbf{MA}(\sigma\text{-centered}) + \neg\mathbf{CH} + \text{“there exist ccc } \Sigma_2^1\text{-notions of forcing } \mathbf{P}_6, \mathbf{P}_6^* \text{ such that } \mathbf{P}_6^* \Vdash \text{“}\hat{\mathbf{P}}_6 \text{ is not ccc”}.$ ■

Problem 7.3 *Is there a ccc Souslin forcing notion $\mathbf{P}$ such that $\mathbf{MA}(\mathbf{P})$ always fails after adding a random real?*

REFERENCES

- [Ba] J.Baumgartner, *Iterated forcing* in Surveys in Set Theory, ed. by A.R.D.Mathias, London Math. Soc., Lecture Notes 87.
- [BJ] J.Bagaria, H.Judah, *Amoeba forcing, Souslin absoluteness and additivity of measure*, Proceedings of the 1989 MSRI Workshop on Set Theory of the Reals (to appear).
- [BaJ] T.Bartoszynski, H.Judah, *Jumping with random reals*,
- [Je] T.Jech, *Set Theory*, Academic Press, New York 1978.
- [Je1] T.Jech, *Multiple Forcing*, Cambridge Tracts in Mathematics 88, Cambridge 1986.
- [JS1] H.Judah, S.Shelah, *Souslin forcing*, Journal of Symbolic Logic, 53(1988).
- [JS2] H.Judah, S.Shelah, *Martin's axioms, measurability and equiconsistency results*, Journal of Symbolic Logic, 54(1989).
- [JR] H.Judah, A.Roslanowski, *On Shelah's amalgamation*, Proceedings of the Winter Institute on Set Theory of the Reals, Bar-Ilan 1991 (to appear).
- [Ra] J.Raisonnier, *A mathematical proof of Shelah's theorem*, Israel Journal of Mathematics 48(1984).
- [Ro] J.Roitman, *Adding a random or a Cohen real: topological consequences and the effect of Martin's axiom*, Fundamenta Mathematicae vol CIII (1979).
- [Sh] S.Shelah, *Can you take Solovay's inaccessible away?*, Israel Journal of Mathematics, 48(1984).
- [Sh1] S.Shelah, *???Vive la Difference I (new version of models with no isomorphic ultra powers*, Proceedings of the Conference in Set Theory, MSRI 10/89,
- [Sh2] S.Shelah, *Proper and Improper Forcing*, in preparation.
- [Ta] M.Talagrand, *Compacts de fonctions mesurables et filtres non mesurables*, Studia Mathematica, 67(1980).
- [To] S.Todorcevic, *Two examples of Borel partially ordered sets with the countable chain conditions*, Proceedings of the American Mathematical Society, 112(1991).